

RELIEF FROM FLIES

Up to 4 weeks
control of:



flies



lice



midges

Deosect[®]

Horses are prone to attack from many species of fly, including the midge *Culicoides*, which are irritating to the horse and can cause disease such as Sweet Itch.

Horses are not only at risk of fly related diseases such as dermatitis, pruritic (itchy) conditions, lumps and skin lesions, they are likely to become stressed and uncomfortable in the summer fly season. Flies can also carry other parasites, which can infect the horse.

Horses can also be affected by lice. Lice typically cause itching of the head, neck and flanks, and usually appear in the winter. This can lead to hair loss and self trauma, and if the infestation is severe it can cause anaemia and make the horse susceptible to other disease.

If a horse has a lice infestation, it is important that all rugs and grooming brushes be treated, as well as other horses which have been in direct contact.

DEOSECT is a product for controlling external parasites. The active ingredient, cypermethrin, is the only prescription product to control flies and midges for up to 4 weeks providing long term relief.

DEOSECT also effectively controls lice.

It can be applied using a sprayer or a sponge to suit your individual horse's needs. With one 250ml tin providing 25 treatments for fly and lice control for your horse, DEOSECT is ready to be prepared, summer or winter – whenever you may need it.

Up to 4 weeks
control of:



flies



lice



midges



250ml = 25 treatments
1 litre = 100 treatments
Up to 4 weeks control

Deosect[®]

Deosect 5% w/v Concentrate for Cutaneous Spray Solution

Presentation

Clear yellow liquid free from solids, containing Cypermethrin technical (cis 50: trans 50) 5.0% w/v (0.1% w/v when diluted).

Uses

Topical spray to control flies and lice on horses.

Dosage and administration

Deosect Spray must be diluted and mixed with water to make a solution just before use. A knapsack, hand sprayer or a sponge may be used to apply the solution. The solution should be used within 24 hours. Avoid spraying directly into animals eyes. One litre (2 pints) of Deosect should be diluted with 50 litres (11 gallons) of water. Alternatively, the following table may be used:

Deosect	Water
10 ml	500 ml
20 ml	1 litre
100 ml	5 litres
200 ml	10 litres

A medium fine spray is advisable to avoid fogging of the diluted Deosect Spray. Careful attention must be paid to total animal coverage, especially the face, legs, back and hind quarters. The following amounts should be used:

Adult horses

500 ml of diluted solution per animal.

Ponies

125 ml of diluted solution per animal.

Fly Control

The application of the solution should commence at the onset of the fly season and be repeated monthly for general fly control and every two weeks for severe head and face fly control.

Lice Control

Apply when lice are observed and repeat as necessary at a minimum interval of 14 days.

Contra-indications, warnings, etc

For external use only.

Do not apply to any lesions where the skin is broken.

Do not spray in or near milking parlours or where milking machinery may be contaminated.

This product belongs to the pyrethroid group. Possible undesirable effects can be irritation and allergic reactions on the skin.

The product can be safely used during pregnancy and lactation.

Deosect spray is of low toxicity and acute poisoning is unlikely to occur.

Operator warnings

FLAMMABLE. Wear impermeable gloves, face shield and rubber boots when handling the concentrate, diluted liquid and when spraying animals.

Use in a well-ventilated area and avoid inhaling the spray mist. If splashed in the eye or on the skin, wash with plenty of clean, running water immediately and seek medical attention.

Remove contaminated clothing immediately and wash before re-use.

Do not smoke or eat while handling the product.

Wash hands and exposed skin after handling this product and before meals.

Withdrawal periods

Not to be used in animals intended for human consumption. Treated horses may never be slaughtered for human consumption.

The horse must have been declared as not intended for human consumption under national horse passport legislation.

Pharmaceutical precautions

Do not store above 25°C. Protect from frost.

Store in original container, tightly closed, in a safe place away from food, drink and animal feedstuffs.

Shelf-life after dilution according to directions: 24 hours.

Dangerous to fish and crustaceans. Do not contaminate ponds, waterways or ditches with the product or used containers.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Keep out of the sight and reach of children. For animal treatment only.

Legal category

POM-VPS

Packaging Quantities

1 litre or 250 ml oblong, unlined tin-plated mild steel can with a tin-plated mild steel cap (screw-fit).

Not all pack sizes may be marketed.

Marketing authorisation number

Vm 42058/4033.

Further information is available from:
Zoetis UK Ltd.,
Walton Oaks, Dorking Road, Walton-on-the-Hill, Tadworth, Surrey KT20 7NS
Customer Support: 0845 3008034
www.zoetis.co.uk

Speak to your medicine prescriber about the use of this or alternative products.

Use medicines responsibly: www.noah.co.uk/responsible/

Further information is available on the SPC

® Registered trademark

Date of preparation: 4.15 AH255/15